Kathmandu University School of Management MPhil Admission Test Sample Question for Quantitative Aptitude Test

Time: 20 minutes/Marks: 20	Symbol no:
----------------------------	------------

(Note: In admission test there are going to be 20 MCQs for 40 marks to do in 40 minutes)

Given below are ten multiple choice questions. *Choose the best answer* for each question by using a circle around your choice. Each question carries equal marks.

- 1. Given $X = \{1,3,5,7,8,9\}$ and $Y = \{2,4,7,9\}$ where X and Y are two sets. Here $X \cap Y$ is:
 - a) {1,3,5,7,8,9, 2,4,7,9}
 - b) {1,3,5,7,8,9,2,4}
 - c) {7,9}
 - d) {1,3,5,8,2,4}
- 2. The results of a survey to 75 adults which asked, "Did you read newspaper last week?" is shown in the table below.

		Did you read newspaper last week		
		Yes	No	
Gender	Male	46	3	
	Female	11	15	

Calculate P (Yes|Male), or the probability that a person responded "Yes" given that the person chosen is male is

- a) 46/49
- b) 46/75
- c) 46/57
- d) 49/75
- 3. What will be the 1st quartile value for the following data set?

- a) 65
- b) 71
- c) 68
- d) 72.5
- 4. One of the following comments on standard deviation is false? Which one?
- a) It is a measure of variation in a data set from its mean.
- b) It has same unit as the unit of the parent data set
- c) Its values are bounded between –ve infinity to +ve infinity
- d) In finance literature, it is used as a measure of risk
- 5. A random variable has binomial distribution with probability of success equal to 0.4. If a sample of size 5 is drawn, then

- a) mean and variance of this variable are 2 and 3 respectively
- b) mean and variance of this variable are 2 and 1.2 respectively
- c) mean and variance of this variable are 5 and 1.2 respectively
- d) mean and variance of the variable can't be calculated from given information
- 6. If a researcher selects every kth item from a population of N items, then she is likely conducting which random sampling?
- a) judgmental
- b) systematic
- c) cluster
- d) stratified
- 7. A researcher is testing a hypothesis on a population mean taking a specific value. The critical z value for α = .05 and a two tailed critical value is \pm 1.96. The observed z value for the sample data is -2.11. The decision made by the researcher based on this information is to
- a) reject the alternative hypothesis
- b) not reject the null hypothesis
- c) increase sample size
- d) reject the null hypothesis
- 8. A researcher finds that in a community 20% of families have average monthly income of Rs 50000 and 80% of families have average monthly income of Rs 20000. Assuming same family size, the overall average monthly income of a family in the community would be
- a) Rs 20000
- b) Rs 26000
- c) Rs 35000
- d) can't be determined from the given information
- 9. Which of the following will most likely cause selection bias in a survey that investigates whether smoking should be prohibited for age under 21?
- a) conducting survey on telephone
- b) receiving only 50% of responses on questionnaire through mail
- c) using of Likert scale rather than having opinion based survey
- d) using a random sample of smokers
- 10. In a regression of Y on X, it was found that: estimated Y = 23.4 + 1.0 X. R^2 value for the regression was 0.64. Which of the following is false?
- a) 64% of variation in Y is explained by the regression
- b) correlation between Y and X is 1
- c) Increase in X by 1 unit increases Y by 1 unit as well
- d) There is a positive relation between Y and X

Answer:

1									
С	a	c	c	b	b	d	b	d	b